

Ryan S. Charette MD Adam Q Ferguson DO Jon Dale PA-C Christa Beyer PA-C Aaron S Covey MD MBA Rania R. Rifaey MD Edward S. Gensicki DPM Megan R. Wolf MD Jon C Driscoll MD Hudson H. Seidel MD Robert C. Stockton DO



www.comcllc.com PH:203.265.3280 FX:203.741.6569

### COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT Brandon Heyda OTR/CHT Raymond Ryan PT/Director Valerie Peckingham PT Isabella Mesturini PT,DPT Peter Ives PTA Kirsten Perillo PTA

# Sports Medicine/Foot & Ankle

## **OLECRANON BURSECTOMY**

Post-Op Protocol

### Phase I—Immediate Motion Phase

Megan R. Wolf, MD

### Goals

- Improve/regain range of motion
- Delay muscular atrophy
- Decrease pain/inflammation

### Days 0-7

- Rest in splint
- Elevate to reduce swelling
- Ice regularly

### Days 7-14

- Range of motion to tolerance (elbow flexion/extension and supination/pronation)
  - o Often full elbow extension is not capable due to pain
- Gentle overpressure into extension
- Wrist flex/ext exercises
- Gripping exercises with putty
- Isometrics for wrist/elbow
- Compression/ice 4-5 times daily

### Days 14-17

- Range of motion ext/flex (at least 20-90 degrees)
- Overpressure into extension (4-5 times daily)
- Joint mobilization to re-establish ROM
- Continue isometrics and gripping exercises
- Continue use of ice

### Days 17-21

- ROM exercises to tolerance (at least 10-100 degrees)
- Overpressure into extension (3-4 times daily)
- Continue joint mobilization techniques
- Initiate light dumbbell programs (PREs)
- Biceps, triceps, wrist flex/ext, sup/pronators
- Continue use of ice post-exercise



Ryan S. Charette MD Adam Q Ferguson DO Jon Dale PA-C Christa Beyer PA-C

Aaron S Covey MD MBA Rania R. Rifaey MD Edward S. Gensicki DPM Megan R. Wolf MD Jon C Driscoll MD Hudson H. Seidel MD Robert C. Stockton DO

Peter Ives PTA

Kirsten Perillo PTA



Post-Op Protocol

**OLECRANON BURSECTOMY** 

www.comcllc.com PH:203.265.3280 FX:203.741.6569

### COMPREHENSIVE PHYSICAL THERAPY

Melinda Amato OTR/L CHT Brandon Heyda OTR/CHT Raymond Ryan PT/Director Valerie Peckingham PT Isabella Mesturini PT,DPT

# . .

# **Sports Medicine/Foot & Ankle**

### Phase II—Intermediate Phase

### Goals

Increase range of motion

Megan R. Wolf, MD

- Improve strength/power/endurance
- Initiate functional activities

### Week 3-4

- Full ROM exercises (4-5 times daily)
- Overpressure into elbow extension
- Continue PRE program for elbow and wrist musculature
- Initiate shoulder program (Thrower's Ten Shoulder Program)
- Continue joint mobilization
- Continue use of ice post-exercise

### Week 4-7

- Continue all exercises listed above
- Initiate light upper body program
- Continue use of ice post-exercise

### Phase III—Advanced Strengthening Program

### Goals

- Improve strength/power/endurance
- Gradual return to functional activities

### **Criteria to Enter Phase III**

- Full non-painful ROM
- No pain or tenderness

### Week 8-12

- Continue PRE program for elbow and wrist
- Continue shoulder program
- Continue stretching for elbow/shoulder
- Initiate interval program and gradually return to sporting activities